

AD-A177 175

A POPULATION OF EXPONENTIALLY DISTRIBUTED INDIVIDUAL
LIFESPANS CANNOT LEA. (U) SOUTH CAROLINA UNIV COLUMBIA
DEPT OF STATISTICS F GUESS ET AL. NOV 86

1/1


UNCLASSIFIED

AFOSR-TR-87-0059 AFOSR-84-0156

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				
1A. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1B. RESTRICTIVE MARKINGS		
2A. SECURITY CLASSIFICATION AUTHORITY FEB 24 1987		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2B. DECLASSIFICATION/DOWNGRADING SCHEDULE		4. PERFORMING ORGANIZATION REPORT NUMBER		
5A. NAME OF PERFORMING ORGANIZATION Department of Statistics		5B. OFFICE SYMBOL (If applicable)		
6A. ADDRESS (City, State and ZIP Code) University of South Carolina Columbia, SC 29208		7A. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research		
8A. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8B. OFFICE SYMBOL (If applicable) NM		
9A. ADDRESS (City, State and ZIP Code) Bolling AFB, DC 20332		9B. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR. 87-0059		
10. SOURCE OF FUNDING NOS.		11. TITLE (Include Security Classification) A Population of Exponentially Distributed Individual Lifespans Cannot Lead to Gompertzian or to Weibull (with Increasing Mortality Rate) Dynamics		
12. PERSONAL AUTHOR(S) Frank Guess and Matthew Witten		13. TYPE OF REPORT Technical		
14. DATE OF REPORT (Yr., Mo., Day) November 1986		15. PAGE COUNT 12		
16. SUPPLEMENTARY NOTATION				
7. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	Gompertzian and Weibull populations; Mixtures of Exponentials Increasing or Decreasing failure (mortality)	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>It is well documented, in the biological literature, that many species throughout the animal kingdom, exhibit Gompertzian or Weibull-like population level survival distributions. Many researchers have long assumed, believed, or otherwise postulated that an individual organism in such a population, survived according to an exponential survival distribution. Using well-known results from reliability theory, it is shown that if every individual in the population is exponentially distributed, then a Gompertzian or Weibull group/population dynamics (or any other dynamics with a strictly increasing mortality rate for some interval) is not possible. This implies that, for species with a population level Gompertzian or Weibull (with the mortality rate strictly increasing) survival curve, some or all of the individual organisms must have nonexponential lifespans.</p>				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22A. NAME OF RESPONSIBLE INDIVIDUAL Maj. Brian W. Woodruff		22B. TELEPHONE NUMBER (Include Area Code) (202) 767-5027		22C. OFFICE SYMBOL NM

DTIC FILE COPY

AD-A177 175

AFOSR-TR- 87-0059

A Population of Exponentially Distributed Individual
Lifespans Cannot Lead to Gompertzian or to
Weibull (with Increasing Mortality Rate) Dynamics

by

Approved for public release;
distribution unlimited.Frank Guess^{*}

Department of Statistics
University of South Carolina
Columbia, SC 29208

and

NCSU Box 8203
Department of Statistics
NC State University
Raleigh, NC 27695-8203

Matthew Witten[†]

[†]Department of Community Health,
School of Medicine
and
Department of Engineering
Mathematics and Computer Science,
Speed Scientific School
University of Louisville
Louisville, KY 40292

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF INTENT TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.
Matthew J. WITTEN
Chief, Technical Information Division

^{*}Work partially supported by the U. S. Air Force Office of Scientific Research under grant number AFOSR-84-0156 and the U. S. Army Research Office under grant number MIPR ARO 139-85 to the University of South Carolina. Currently on leave of absence from NC State University.

[†]Address all correspondence to this author at this address (+).

Guess and WittenABSTRACT

It is well documented, in the biological literature, that many species throughout the animal kingdom, exhibit Gompertzian or Weibull-like population level survival distributions. Many researchers have long assumed, believed, or otherwise postulated that an individual organism, in such a population, survived according to an exponential survival distribution. Using well-known results from reliability theory, it is shown that if every individual in the population is exponentially distributed, then a Gompertzian or Weibull group/population dynamics (or any other dynamics with a strictly increasing mortality rate for some interval) is not possible. This implies that, for species with a population level Gompertzian or Weibull (with the mortality rate strictly increasing) survival curve, some or all of the individual organisms must have nonexponential lifespans.

Accession For	
NTIS CRAXI	N
DTIC TAB	[]
Unannounced	[]
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Guess and Witten

The biological literature is replete with illustrations of various species of organisms that exhibit Gompertzian, Weibull or Gompertzian like survival distributions (Figures 1-2, Comfort (1), Witten (2-4) for examples). The assumption has long been held, among researchers, that an individual organism-in a population-survives according to an exponential survival distribution. That is, if $S(t)$ is the probability that an individual survives past time t , then $S(t)$ is of the form

$$S(t) = e^{-\lambda t} \quad (1)$$

where λ is the inverse of the expected lifespan. Each individual, in the population could, of course, have a different mortality rate (5).

Exponential survival distributions are derived from the assumption of constant mortality rate (failure rate). This assumption is equivalent to the statement that the probability of an organism's survival until time $t + \Delta t$, given its survival until time t , is constant for all time t . That is, the per capita death rate λ given by

$$-\frac{1}{N} \frac{dN}{dt} = \lambda \quad (2)$$

is a constant. Given the known classes of experimental survival curves, it is useful and important to address the question of whether or not one can derive a Gompertzian lifespan distribution from some mixture (population) of individuals having exponential survival (lifespan) curves. That is, can a population of exponential survival distributions, at the individual level, give rise to a Gompertzian or Weibull-like population dynamics?

Proschan (7) considered a naturally occurring mixture of exponentials in the aircraft industry (Cf. also, for example, (8), (9), (10) and (11) for more on mixtures). Proschan demonstrated that a survival distribution

Guess and Witten

arising from any mixture of exponentials will always possess a decreasing mortality rate (DMR). That is, it will process a decreasing failure rate (where by decreasing we mean nonincreasing). The Weibull-like populations that have been empirically observed are observed to have a strictly increasing mortality rate. Gompertzian populations always have a strictly increasing mortality rate. Given the result of Proschan (7), it is therefore not possible to obtain a Gompertzian or Weibull-like population dynamics from any mixture of individual exponential lifespans; as such a survival distribution implies the existence of a strictly increasing mortality rate. (It is possible, however, for a mixture of exponentials to yield a Weibull with a DMR, e.g., see (10), for neonates a DMR is plausible, but for older organisms it is not.)

The biological import of this statement is that some or all of the individual lifespans, comprising the population, must be nonexponential in their survival behavior. This result will also be the case, more generally, for any population survival distribution with a strictly increasing mortality rate over the entire lifespan or, more restrictively, even only on some subinterval of the lifespan (e.g., for all ages t such that $t \geq t_0$ for t_0 a large enough age). A simple illustration of such a population would be in which there was both neonatal and post-neonatal death (Witten (4)). Here, one mixes exponential and Gompertzian distributions to describe the known population dynamics (Figures 3-4). That is, again some or all individual lifespans will have to be nonexponentially distributed to yield the empirically observed population dynamics.

Guess and WittenReferences and Notes

1. A. Comfort, The Biology of Senescence (Elsevier/North Holland, NY, 1979).
2. M. Witten, Mech. Aging and Dev. 27, 323 (1984).
3. M. Witten, Mech. Aging and Dev. 32, 141 (1985).
4. M. Witten, Mech. Aging and Dev. 33, 177 (1986).
5. One possible means of addressing this issue is to assume that there is some type of distribution $\lambda(t)$ of mortality rates such that $E[\lambda] = \lambda_D(t)$ a deterministic function and that $\lambda(t)$ is perturbed by a mean zero, variance σ^2 , white noise (or colored noise) process. This gives rise to a stochastic differential equation for the survival distribution $S(t)$ (see Witten (6) for discussion and further references).
6. M. Witten, Bull. Math. Biol. 45, 425 (1983).
7. F. Proschan, Technometrics 5, 375 (1963).
8. R. E. Barlow, A. W. Marshall, and F. Proschan, Ann. Math. Stat. 34, 375 (1963).
9. R. Barlow and F. Proschan, Statistical Theory of Reliability and Life Testing (Hold, Rinehart, and Winston, New York, 1975).
10. N. Jewell, Annals of Statistics 10, 479 (1982).
11. M. Hollander and F. Proschan, in Handbook of Statistics, Vol. 4, Nonparametric Methods, P. R. Krishnaiah and P. K. Sen, Eds. (North-Holland, Amsterdam, 1984), pp. 613-655.
12. We gratefully thank F. Proschan for introducing us to each others work. F. G. expresses appreciation to F. Proschan and the Reliability Center at

Guess and Witten

Florida State University for introducing him to general reliability operations on classes of survival distributions. M. W. would like to thank C. E. Finch and the participants of the recent Brookhaven Biology of Aging Conference for their many fruitful discussions.

Guess and Witten

LEGENDS FOR FIGURES

Figure 1: Percent survival of *Cynolebias adoloffi*. Reproduced from Comfort (1).

Figure 2: The effects of NDGA-feeding on survival, in male and female mosquitos. Reproduced courtesy of C. Lang.

Figure 3: Survival curves for United States females in the years 1900, 1960, and 1980. Notice the early neonatal dip in all of these curves. Reproduced courtesy of the author K. Manton.

Guess and Witten

Figure 4: An illustration of the wide variety of dynamics embedded in a two group population model. Reproduced from Witten (4).





Figure 1

Guess and Witten




Guess and Witten

Figure 3

Guess and Witten




Figure 4

END

3-87

DTIC